



Information for Finding a Certified Treatment Center

Indication

SPRAVATO™ (esketamine) CIII Nasal Spray is indicated, in conjunction with an oral antidepressant (AD), for the treatment of treatment-resistant depression (TRD) in adults.

SPRAVATO™ is not approved as an anesthetic agent. The safety and effectiveness of SPRAVATO™ as an anesthetic agent have not been established.

Important Safety Information

WARNING: SEDATION, DISSOCIATION; ABUSE AND MISUSE; and SUICIDAL THOUGHTS AND BEHAVIORS

See full prescribing information for complete boxed warning


- **Risk for sedation and dissociation after administration. Monitor patients for at least two hours after administration (5.1, 5.2).**
- **Potential for abuse and misuse. Consider the risks and benefits of using SPRAVATO™ prior to use in patients at higher risk of abuse. Monitor for signs and symptoms of abuse and misuse (5.3).**
- **SPRAVATO™ is only available through a restricted program called the SPRAVATO™ REMS (5.4).**
- **Increased risk of suicidal thoughts and behaviors in pediatric and young adult patients taking antidepressants. Closely monitor all antidepressant-treated patients for clinical worsening and emergence of suicidal thoughts and behaviors. SPRAVATO™ is not approved for use in pediatric patients (5.5).**

Please see Important Safety Information, including Boxed WARNINGS, on pages 4-7.

Please see pocketed full Prescribing Information, including Boxed WARNINGS, for SPRAVATO™.

Information for Finding a Certified Treatment Center

Once a patient is ready to begin treatment with SPRAVATO™ (esketamine) CIII Nasal Spray, they can start treatment in a certified SPRAVATO™ treatment center. SPRAVATO™ must be administered under the direct supervision of a healthcare provider.¹ Certified SPRAVATO™ treatment centers will provide the infrastructure and operational support to prepare and monitor patients during self-administration and prior to release from the treatment center. If you choose to refer patients to a certified SPRAVATO™ treatment center, it may be helpful to share complete and timely patient information among providers for continued patient care.



SEEKING NEW TREATMENT

Discuss SPRAVATO™ Treatment

- Discuss SPRAVATO™ in conjunction with an oral antidepressant (AD) as a treatment option for appropriate adult patients with treatment-resistant depression (TRD)
- Communicate the benefits and risks associated with SPRAVATO™ to patients and caregivers

Select a Certified SPRAVATO™ Treatment Center

- Identify a certified SPRAVATO™ treatment center that meets your patient's needs
- It is important to share patient- and practice-related information with the certified SPRAVATO™ treatment center for quality consultation and coordinated patient care
- You may use your current practice protocol for referrals or see the next page for information for your consideration to help facilitate effective referrals

For more information about how to find a certified SPRAVATO™ treatment center near you, visit www.spravatohcp.com/findacenter or call a Janssen CarePath Care Coordinator at 844-777-2828, Monday-Friday, 8:00 AM to 8:00 PM ET.

Share Next Steps

- Referring physicians or patients may contact a certified SPRAVATO™ treatment center for additional information and to schedule an initial consultation for SPRAVATO™ treatment

Communicating patient information at the time of referral supports quality consultation and coordinated patient care among providers. It is estimated that timely referral communication results in a 2-fold increase in patient access to optimal care. Sharing incomplete information may cause delays in patient access to treatment and increase administrative time.²

For additional information about how to become a treatment center, please visit www.spravatotreatmentcenter.com.

**Please see Important Safety Information, including Boxed WARNINGS, on pages 4-7.
Please see pocketed full Prescribing Information, including Boxed WARNINGS, for SPRAVATO™.**

Considerations to Help Facilitate Effective Referrals

Connect your patients to a certified SPRAVATO™ treatment center

Sharing patient information with a certified SPRAVATO™ treatment center may help ensure timely coordination of care. You or your patients may choose to reference these considerations to share patient- and practice-related information with a certified SPRAVATO™ treatment center.



Patient Information

- Name
- Date of Birth
- Gender
- Patient's Phone Number
- Address
- Caregiver Name and Phone Number



Patient Insurance Information

- Primary Insurance
- Policy #
- Group #
- Policyholder's Name
- Card/BIN #



Certified SPRAVATO™ Treatment Center Information

- Name
- Address
- Name and Phone Number
- Email



Referring Physician's Information

- Name
- Address
- Name and Phone Number
- Email



Clinical Information

- Diagnosis
- Laboratory Procedure/Tests and Dates
- Medical History
- Medication History

janssen
CarePath

Janssen CarePath helps verify insurance coverage for patients, provides reimbursement information, helps find financial assistance options for eligible patients, and provides ongoing support to help patients start and stay on prescribed SPRAVATO™ treatment.

Call a Janssen CarePath Care Coordinator at 844-777-2828,
Monday-Friday, 8:00 AM to 8:00 PM ET.

Visit JanssenCarePath.com/hcp/Spravato.

Spravato™
(esketamine) III
nasal spray

Important Safety Information for SPRAVATO™ (esketamine) CIII Nasal Spray

Indication

SPRAVATO™ (esketamine) CIII Nasal Spray is indicated, in conjunction with an oral antidepressant (AD), for the treatment of treatment-resistant depression (TRD) in adults.

SPRAVATO™ is not approved as an anesthetic agent. The safety and effectiveness of SPRAVATO™ as an anesthetic agent have not been established.

Important Safety Information

WARNING: SEDATION, DISSOCIATION; ABUSE AND MISUSE; and SUICIDAL THOUGHTS AND BEHAVIORS

See full prescribing information for complete boxed warning

- **Risk for sedation and dissociation after administration. Monitor patients for at least two hours after administration (5.1, 5.2).**
- **Potential for abuse and misuse. Consider the risks and benefits of using SPRAVATO™ prior to use in patients at higher risk of abuse. Monitor for signs and symptoms of abuse and misuse (5.3).**
- **SPRAVATO™ is only available through a restricted program called the SPRAVATO™ REMS (5.4).**
- **Increased risk of suicidal thoughts and behaviors in pediatric and young adult patients taking antidepressants. Closely monitor all antidepressant-treated patients for clinical worsening and emergence of suicidal thoughts and behaviors. SPRAVATO™ is not approved for use in pediatric patients (5.5).**

CONTRAINDICATIONS

SPRAVATO™ is contraindicated in patients with:

- Aneurysmal vascular disease (including thoracic and abdominal aorta, intracranial and peripheral arterial vessels) or arteriovenous malformation
- History of intracerebral hemorrhage
- Hypersensitivity to esketamine, ketamine, or any of the excipients

WARNINGS AND PRECAUTIONS

Sedation: In clinical trials, 49% to 61% of SPRAVATO™-treated patients developed sedation and 0.3% of SPRAVATO™-treated patients experienced loss of consciousness.

Because of the possibility of delayed or prolonged sedation, patients must be monitored by a healthcare provider for at least 2 hours at each treatment session, followed by an assessment to determine when the patient is considered clinically stable and ready to leave the healthcare setting.

Closely monitor for sedation with concomitant use of SPRAVATO™ with CNS depressants [see *Drug Interaction (7.1)*]. SPRAVATO™ is available only through a restricted program under a REMS.

Dissociation: The most common psychological effects of SPRAVATO™ were dissociative or perceptual changes (including distortion of time, space and illusions), derealization and depersonalization (61% to 75% of SPRAVATO™-treated patients developed dissociative or perceptual changes). Given its potential to induce dissociative effects, carefully assess patients with psychosis before administering SPRAVATO™; treatment should be initiated only if the benefit outweighs the risk.

Because of the risks of dissociation, patients must be monitored by a healthcare provider for at least 2 hours at each treatment session, followed by an assessment to determine when the patient is considered clinically stable and ready to leave the healthcare setting.

SPRAVATO™ is available only through a restricted program under a REMS.

Abuse and Misuse: SPRAVATO™ contains esketamine, a Schedule III controlled substance (CIII), and may be subject to abuse and diversion. Assess each patient's risk for abuse or misuse prior to prescribing and monitor all patients for the development of these behaviors or conditions, including drug-seeking behavior, while on therapy. Individuals with a history of drug abuse or dependence are at greater risk; therefore, use careful consideration prior to treatment of individuals with a history of substance use disorder and monitor for signs of abuse or dependence.

SPRAVATO™ is available only through a restricted program under a REMS.

SPRAVATO™ Risk Evaluation and Mitigation Strategy (REMS):

SPRAVATO™ is available only through a restricted program called the SPRAVATO™ REMS because of the risks of serious adverse outcomes from sedation, dissociation, and abuse and misuse.

Important requirements of the SPRAVATO™ REMS include the following:

- Healthcare settings must be certified in the program and ensure that SPRAVATO™ is:
 - Only dispensed in healthcare settings and administered to patients who are enrolled in the program.
 - Administered by patients under the direct observation of a healthcare provider and that patients are monitored by a healthcare provider for at least 2 hours after administration of SPRAVATO™.
- Pharmacies must be certified in the REMS and must only dispense SPRAVATO™ to healthcare settings that are certified in the program.

Please see pocketed full Prescribing Information, including Boxed WARNINGS, for SPRAVATO™.

Further information, including a list of certified pharmacies, is available at www.SPRAVATOrems.com or 1-855-382-6022.

Suicidal Thoughts and Behaviors in Adolescents and Young Adults:

In pooled analyses of placebo-controlled trials of antidepressant drugs (SSRIs and other antidepressant classes) that included adult and pediatric patients, the incidence of suicidal thoughts and behaviors in patients age 24 years and younger was greater than in placebo-treated patients. SPRAVATO™ is not approved in pediatric (<18 years of age) patients.

There was considerable variation in risk of suicidal thoughts and behaviors among drugs, but there was an increased risk identified in young patients for most drugs studied.

Monitor all antidepressant-treated patients for clinical worsening and emergence of suicidal thoughts and behaviors, especially during the initial few months of drug therapy and at times of dosage changes. Counsel family members or caregivers of patients to monitor for changes in behavior and to alert the healthcare provider. Consider changing the therapeutic regimen, including possibly discontinuing SPRAVATO™ and/or the concomitant oral antidepressant, in patients whose depression is persistently worse, or who are experiencing emergent suicidal thoughts or behaviors.

Increase in Blood Pressure: SPRAVATO™ causes increases in systolic and/or diastolic blood pressure (BP) at all recommended dosages. Increases in BP peak approximately 40 minutes after SPRAVATO™ administration and last approximately 4 hours.

Approximately 8% to 17% of SPRAVATO™-treated patients experienced an increase of more than 40 mmHg in systolic BP and/or 25 mmHg in diastolic BP in the first 1.5 hours after administration at least once during the first 4 weeks of treatment. A substantial increase in blood pressure could occur after any dose administered even if smaller blood pressure effects were observed with previous administrations. SPRAVATO™ is contraindicated in patients for whom an increase in BP or intracranial pressure poses a serious risk (e.g., aneurysmal vascular disease, arteriovenous malformation, history of intracerebral hemorrhage). Before prescribing, patients with other cardiovascular and cerebrovascular conditions should be carefully assessed to determine whether the potential benefits of SPRAVATO™ outweigh its risk.

Assess BP prior to administration of SPRAVATO™. In patients whose BP is elevated prior to SPRAVATO™ administration (as a general guide: >140/90 mmHg), a decision to delay SPRAVATO™ therapy should be taken into account to balance the benefit and risk in individual patients.

BP should be monitored for at least 2 hours after SPRAVATO™ administration. Measure blood pressure around 40 minutes post-dose and subsequently as clinically warranted until values decline. If BP remains high, promptly seek assistance from practitioners experienced in BP management. Refer patients experiencing symptoms of a hypertensive crisis (e.g., chest pain, shortness of breath) or hypertensive encephalopathy (e.g., sudden severe headache, visual disturbances, seizures, diminished consciousness or focal neurological deficits) immediately for emergency care.

Closely monitor blood pressure with concomitant use of SPRAVATO™ with psychostimulants or monoamine oxidase inhibitors (MAOIs) [see *Drug Interactions (7.2, 7.3)*].

In patients with history of hypertensive encephalopathy, more intensive monitoring, including more frequent blood pressure and symptom assessment, is warranted because these patients are at increased risk for developing encephalopathy with even small increases in blood pressure.

Cognitive Impairment

Short-Term Cognitive Impairment: In a study in healthy volunteers, a single dose of SPRAVATO™ caused cognitive performance decline 40 minutes post-dose. SPRAVATO™-treated subjects required a greater effort to complete the cognitive tests at 40 minutes post-dose. Cognitive performance and mental effort were comparable between SPRAVATO™ and placebo at 2 hours post-dose. Sleepiness was comparable after 4 hours post-dose.

Long-Term Cognitive Impairment: Long-term cognitive and memory impairment have been reported with repeated ketamine misuse or abuse. No adverse effects of SPRAVATO™ nasal spray on cognitive functioning were observed in a one-year open-label safety study; however, the long-term cognitive effects of SPRAVATO™ have not been evaluated beyond one year.

Impaired Ability to Drive and Operate Machinery:

Before SPRAVATO™ administration, instruct patients not to engage in potentially hazardous activities requiring complete mental alertness and motor coordination, such as driving a motor vehicle or operating machinery, until the next day following a restful sleep. Patients will need to arrange transportation home following treatment with SPRAVATO™.

(continued on next page)



Important Safety Information for SPRAVATO™ (esketamine) CIII Nasal Spray (continued)

Ulcerative or Interstitial Cystitis: Cases of ulcerative or interstitial cystitis have been reported in individuals with long-term off-label use or misuse/abuse of ketamine. In clinical studies with SPRAVATO™ nasal spray, there was a higher rate of lower urinary tract symptoms (pollakiuria, dysuria, micturition urgency, nocturia, and cystitis) in SPRAVATO™-treated patients than in placebo-treated patients. No cases of esketamine-related interstitial cystitis were observed in any of the studies, which involved treatment for up to a year.

Monitor for urinary tract and bladder symptoms during the course of treatment with SPRAVATO™ and refer to an appropriate healthcare provider as clinically warranted.

Embryo-fetal Toxicity: SPRAVATO™ may cause fetal harm when administered to pregnant women. Advise pregnant women of the potential risk to an infant exposed to SPRAVATO™ *in utero*. Advise women of reproductive potential to consider pregnancy planning and prevention.

DRUG INTERACTIONS

CNS depressants (e.g., benzodiazepines, opioids, alcohol): Concomitant use may increase sedation. Closely monitor for sedation with concomitant use of CNS depressants.

Psychostimulants (e.g., amphetamines, methylphenidate, modafinil, armodafinil): Concomitant use may increase blood pressure. Closely monitor blood pressure with concomitant use of psychostimulants.

Monoamine oxidase inhibitors (MAOIs): Concomitant use may increase blood pressure. Closely monitor blood pressure with concomitant use of MAOIs.

USE IN SPECIFIC POPULATIONS

Pregnancy: SPRAVATO™ is not recommended during pregnancy. SPRAVATO™ may cause fetal harm when administered to pregnant women. Advise pregnant women of the potential risk to an infant exposed to SPRAVATO™ *in utero*. There are risks to the mother associated with untreated depression in pregnancy. If a woman becomes pregnant while being treated with SPRAVATO™, treatment with SPRAVATO™ should be discontinued and the patient should be counseled about the potential risk to the fetus.

Pregnancy Exposure Registry: There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to antidepressants, including SPRAVATO™, during pregnancy. Healthcare providers are encouraged to register patients by contacting the National Pregnancy Registry for Antidepressants at 1-844-405-6185 or online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/>.

Lactation: SPRAVATO™ is present in human milk. Because of the potential for neurotoxicity, advise patients that breastfeeding is not recommended during treatment with SPRAVATO™.

Females and Males of Reproductive Potential: SPRAVATO™ may cause embryo-fetal harm when administered to a pregnant woman. Consider pregnancy planning and prevention for females of reproductive potential during treatment with SPRAVATO™.

Pediatric Use: The safety and effectiveness of SPRAVATO™ in pediatric patients have not been established.

Geriatric Use: Of the total number of patients in Phase 3 clinical studies exposed to SPRAVATO™, 12% were 65 years of age and older, and 2% were 75 years of age and older. No overall differences in the safety profile were observed between patients 65 years of age and older and patients younger than 65 years of age.

The mean esketamine C_{max} and AUC values were higher in elderly patients compared with younger adult patients.

The treatment of TRD in geriatric patients was evaluated in a 4-week, randomized, double-blind study comparing flexibly-dosed intranasal SPRAVATO™ plus a newly initiated oral antidepressant compared to intranasal placebo plus a newly initiated oral antidepressant in patients ≥ 65 years of age. At the end of four weeks, there was no statistically significant difference between groups on the primary efficacy endpoint of change from baseline to Week 4 on the Montgomery-Åsberg Depression Rating Scale (MADRS).

Hepatic Impairment: SPRAVATO™-treated patients with moderate hepatic impairment may need to be monitored for adverse reactions for a longer period of time.

SPRAVATO™ has not been studied in patients with severe hepatic impairment (Child-Pugh class C). Use in this population is not recommended.

DRUG ABUSE AND DEPENDENCE

Controlled Substance: SPRAVATO™ contains esketamine hydrochloride, the (S)-enantiomer of ketamine and a Schedule III controlled substance under the Controlled Substances Act.

Abuse: Individuals with a history of drug abuse or dependence may be at greater risk for abuse and misuse of SPRAVATO™. Abuse is the intentional, non-therapeutic use of a drug, even once, for its psychological or physiological effects. Misuse is the intentional use, for therapeutic purposes, of a drug by an individual in a way other than prescribed by a healthcare provider or for whom it was not prescribed. Careful consideration is advised prior to use of individuals with a history of substance use disorder, including alcohol.

Please see pocketed full Prescribing Information, including Boxed WARNINGS, for SPRAVATO™.

SPRAVATO™ may produce a variety of symptoms including anxiety, dysphoria, disorientation, insomnia, flashback, hallucinations, and feelings of floating, detachment and to be “spaced out.” Monitoring for signs of abuse and misuse is recommended.

ADVERSE REACTIONS

The most common adverse reactions with SPRAVATO™ plus oral AD (incidence ≥5% and at least twice that of placebo nasal spray plus oral AD) were dissociation, dizziness, nausea, sedation, vertigo, hypoesthesia, anxiety, lethargy, blood pressure increased, vomiting, and feeling drunk.

Please see pocketed full Prescribing Information, including Boxed WARNINGS, for SPRAVATO™.

cp-79821v1

Spravato™
(esketamine) Ⓒ
nasal spray 